

National Pest Alert

Zika Virus

Zika Virus Transmission

Zika, a mosquito-transmitted virus, has recently been reported in North America from travelers who have returned from areas where the virus is present. The primary means of transmission is from the bite of infected female *Aedes* mosquitoes. Other modes of transmission that have been documented include blood transfusion, sexual contact, and rarely, from mother to child.

As of August 2016, local transmission of the Zika virus has been reported in the United States via mosquitoes as well as imported and locally transmitted through sexual transmission from an infected partner. Local transmission of the virus is occurring in the following states and territories of the United States: Florida, American Samoa, Puerto Rico, Marshall Islands, and the US Virgin Islands (St. Croix, St. Thomas, and St. John). For the most current information on local transmission of the virus visit wwwnc.cdc.gov/travel/page/zika-travel-information. The primary mosquito vector appears to be the yellow fever mosquito, *Aedes aegypti*, although a number of other *Aedes* species are known to serve as vectors capable of transmitting the disease to humans including the Asian tiger mosquito (*Aedes albopictus*).

Origin of Zika

Zika was first isolated in 1947 in Uganda's Zika Forest in Africa where it was discovered in a rhesus monkey. The virus remained relatively unknown until a large outbreak occurred in Micronesia in 2007. Scientists believe the Zika virus mutated into the 2007 increased virulence strain from the original strain found in Uganda. Since 2007 the virus has rapidly spread through the Pacific Islands and in 2015 it appeared in South America.

Mosquito Life Cycle

Mosquitoes are classified into the order Diptera (true flies) and have four distinct life stages (egg, larva, pupa, and adult). Female mosquito species known to transmit Zika virus lay their eggs on moist surfaces such as the interior walls of treeholes, cans, bromeliad plants, various types of containers, and old tires that are likely to be flooded by water. Most larvae hatch within 48 hours and the larvae and pupae live in water. The adult mosquito emerges from the pupal case and rests on the water's surface until its body dries




Ed Freytag

 *Aedes aegypti*

 *Aedes albopictus*

 both *A. aegypti* and *A. albopictus*



Known distribution is estimated and actual distribution may shift quickly due to mosquito movement and weather patterns.

and its exoskeleton hardens. Male and female mosquitoes generally feed on flower nectar. Female mosquitoes require a blood meal before they can lay eggs, so only female mosquitoes bite. They bite every few days during their entire adult lives, which may last several weeks.

Mosquito Prevention and Control

You can most effectively reduce the number of mosquitoes around your home and neighborhood by eliminating the standing



United States Department of Agriculture

National Institute of Food and Agriculture



water in which mosquitoes require to complete their development cycle.

- Dispose of any refuse that can hold water, such as tin cans, containers, and in particular, used tires. Tires have become the most important developmental sites for Zika-transmitting mosquitoes in the U.S. Even small containers holding less than one cup of water are sufficient for the development of these mosquitoes.
- Drill holes in the bottoms of recycling containers and check uncovered junk piles.
- Clean clogged roof gutters every year, and check storm drains, leaky outdoor faucets, and window wells. Fallen leaves and other organic debris indirectly provide food for mosquito larvae.
- Prevent the accumulation of standing water and empty water from wheelbarrows, boats, cargo trailers, pet dishes, toys, saucers underneath flower pots, and ceramic pots. If possible, turn these items over when not in use.
- Do not allow water to stagnate in birdbaths, ornamental pools, water gardens, and swimming pools or their covers. Ornamental pools can be aerated or stocked with fish. Swimming pools should be cleaned and chlorinated when not in use.
- Alter the landscape of your property to eliminate standing water. *Keep in mind that during warm weather, mosquitoes can develop in any puddle of water.* Larvicides are highly effective in controlling immature mosquitoes and should be considered when standing water cannot be eliminated.

Protect Yourself from Bites

Even though your property may lack mosquito-developmental sites, mosquitoes can travel from their aquatic habitat site in search of a blood meal. Therefore, it may be necessary to supplement larval control with other control measures directed at adult mosquitoes. The following tips can help to reduce your risk of being bitten by a mosquito:

- Make sure window and door screens are “bug tight.” Window screening is made to keep out mosquitoes, so repair any holes or tears. Install weather-stripping to keep mosquitoes from entering around loose fitting doors and windows.
- Use the proper type of light outside: incandescent lights attract mosquitoes, whereas fluorescent lights neither attract nor repel mosquitoes.
- *Aedes* species associated with the Zika virus are considered “day” biters so protect yourself whenever outside or in the presence of mosquitoes. Long-sleeved shirts and long pants provide some protection against bites.
- Insect repellents when applied according to the label directions to exposed skin deter mosquitoes from biting. Spray thin clothing with repellent because mosquitoes can bite through it. Various repellents differ in how they are formulated. Some are labelled for application to skin only, some can be applied to clothing only, and some are labeled for both skin and clothing. It is important to read the label of the product to determine how it should be applied. The American Academy of Pediatrics recommends that

repellents used on children contain no more than 10 percent DEET, the active ingredient in mosquito repellents. The Centers for Disease Control guidelines state that pregnant women are not excluded from using EPA-registered repellents. Be sure to follow all directions on product labels.

Symptoms of the Zika Virus

Most humans infected with Zika have very mild or no symptoms so many individuals may not realize they are infected with the virus. A small number of infected individuals (1 in 5) develop mild symptoms that include fever, joint pain, conjunctivitis and a body rash. Symptoms typically occur 2 to 7 days following the bite from an infected mosquito. More severe symptoms may occur in some individuals including paralysis. Researchers are investigating the link between the Zika virus and birth defects including microcephaly.

The World Health Organization and the Centers for Disease Control and Prevention are tracking the distribution of the virus and recommending some individuals including pregnant women refrain from traveling to areas where the virus is present. State health departments and university extension personnel may have mosquito control and detection programs for your state.

For more information about the Zika virus, mosquito control recommendations, and state resources visit our Web site at: ncipmc.org/action/alerts/zika.php

This publication was produced and distributed in cooperation with the USDA NIFA Regional Integrated Pest Management Program Centers and State-based Extension Programs. For more information regarding the development of this document, please contact Susan T. Ratcliffe at sratclif@illinois.edu or by phone at (217) 333-9656.

Contributors:

Nathan Burkett-Cadena, University of Florida
C. Roxanne Connelly, University of Florida
Janet Hurley, Texas A&M University
Phil Nixon, University of Illinois
Faith Oi, University of Florida
Susan Ratcliffe, University of Illinois

Aedes images from:

Ed Freytag, New Orleans Mosquito, Termite and Rodent Control Board

1862 Land-Grant Universities

Auburn University University of Alaska University of Arizona University of Arkansas University of California Colorado State University University of Connecticut University of Delaware University of the District of Columbia University of Florida University of Georgia University of Guam University of Hawaii University of Idaho University of Illinois Purdue University Iowa State University Kansas State University University of Kentucky Louisiana State University University of Maine University of Maryland University of Massachusetts Michigan State University University of Minnesota Mississippi State University University of Missouri Montana State University University of Nebraska University of Nevada University of New Hampshire Rutgers New Mexico State University Cornell University North Carolina State University North Dakota State University Ohio State University Oklahoma State University Oregon State University Pennsylvania State University University of Puerto Rico University of Rhode Island Clemson University South Dakota State University University of Tennessee Texas A&M University Utah State University University of Vermont University of the Virgin Islands Virginia Polytechnic Institute & State University Washington State University West Virginia University West Virginia State University University of Wisconsin University of Wyoming.

August 2016



United States Department of Agriculture National Institute of Food and Agriculture

